

US FDA Type C Meeting for ATL1102 in DMD

Antisense Therapeutics Limited [ASX:ANP | US OTC:ATHJY | FSE:AWY], (the Company) is pleased to advise that Type C guidance meeting with the US Food and Drug Administration (FDA) was held as scheduled on 19 April 2021 ET to discuss the further development of ATL1102 in Duchenne muscular dystrophy (DMD) in the US. The Company expects to provide further details following receipt from the FDA of the official minutes of the meeting in late May 2021.

The meeting was constructive and provided clarification on a path towards initiating a Phase IIb/III study in the US. Non-clinical requirements are to be further reviewed and agreed with the FDA to assess if and how they may impact on the timing of clinical study initiation. The feedback will be reviewed by our internal team and incorporated into our global clinical development and commercialisation plans.

Mark Diamond CEO of Antisense Therapeutics said: "Based on the guidance meeting with the FDA, together with prior feedback from EMA and world leading DMD experts, we are reassured that the data from our Phase II study is encouraging and that it is reasonable and appropriate to advance the program towards potentially pivotal clinical studies. It is our goal to bring this medication to as many DMD patients worldwide as possible".

This announcement has been authorised for release by the Board

For more information please contact:

Antisense Therapeutics

Mark Diamond
Managing Director
+61 (0)3 9827 8999
www.antisense.com.au

Investment Enquiries

Gennadi Koutchin
XEC Partners
gkoutchin@xecpartners.com.au
1300 932 037

About Antisense Therapeutics Limited [ASX: ANP | US OTC: ATHJY | FSE: AWY] is an Australian publicly listed biotechnology company, developing and commercializing antisense pharmaceuticals for large unmet markets in rare diseases. The products are in-licensed from Ionis Pharmaceuticals Inc. (NASDAQ: IONS), an established leader in antisense drug development. The Company is developing ATL1102, an antisense inhibitor of the CD49d receptor, for Duchenne muscular dystrophy (DMD) patients and recently reported highly promising Phase II trial results. ATL1102 has also successfully completed a Phase II efficacy and safety trial, significantly reducing the number of brain lesions in patients with relapsing-remitting multiple sclerosis (RRMS). The Company has a second drug, ATL1103 designed to block GHr production that successfully reduced blood IGF-I levels in Phase II clinical trials in patients with the growth disorder acromegaly.

About ATL1102 ATL1102 is an antisense inhibitor of CD49d, a subunit of VLA-4 (Very Late Antigen-4). Antisense inhibition of VLA-4 expression has demonstrated activity in a number of animal models of inflammatory disease including asthma and MS with the MS animal data having been published in a peer reviewed scientific journal. ATL1102 was shown to be highly effective in reducing MS lesions in a Phase IIa clinical trial in patients with RR-

MS. The ATL1102 Phase IIa clinical data has been published in the medical Journal **Neurology** (Limmroth, V. et al Neurology, 2014; 83(20): 1780-1788).

About DMD Duchenne Muscular Dystrophy (DMD) is an X-linked disease that affects 1 in 3600 to 6000 live male births (Bushby *et al*, 2010). DMD occurs as a result of mutations in the dystrophin gene which causes a substantial reduction in or absence of the dystrophin protein. Children with DMD have dystrophin deficient muscles and are susceptible to contraction induced injury to muscle that triggers the immune system which exacerbates muscle damage as summarized in a publication co-authored by the Director of the FDA CDER (Rosenberg et al, 2015). Ongoing deterioration in muscle strength affects lower limbs leading to impaired mobility, and also affects upper limbs, leading to further loss of function and self-care ability. The need for wheelchair use can occur in early teenage years for patients on corticosteroids with a mean age of 13, with respiratory, cardiac, cognitive dysfunction also emerging. Patients with a greater number of immune T cells expressing high levels of CD49d have more severe and progressive disease and are non-ambulant by the age of 10 despite being on corticosteroid treatment (Pinto Mariz et al, 2015). With no intervention, the mean age of life is approximately 19 years. The management of the inflammation associated with DMD is currently addressed via the use of corticosteroids, however they are acknowledged as providing insufficient efficacy and are associated with significant side effects. As a consequence, there is an acknowledged high need for new therapeutic approaches for the treatment of inflammation associated with DMD.

Rosenberg AS, Puig M, Nagaraju K, *et al*. Immune-mediated pathology in Duchenne muscular dystrophy. *Sci Transl Med* 2015, 7: 299rv4.

Bushby et al for the DMD Care Consideration Working Group/ *Diagnosis and management of Duchenne muscular dystrophy, part 1* Lancet Neurol. **2010** Jan;9(1):77-93 *and part 2* Lancet Neurol. **2010** Feb;9(2):177-89 .

Pinto-Mariz F, Carvalho LR, Araújo AQC, *et al*. CD49d is a disease progression biomarker and a potential target for immunotherapy in Duchenne muscular dystrophy. *Skeletal Muscle* 2015, 5: 45-55.