
UNITED STATES
SECURITIES AND EXCHANGE COMMISSION
Washington, D.C. 20549

FORM 6-K

Report of Foreign Private Issuer
Pursuant to Rule 13a-16 or 15d-16
of the Securities Exchange Act of 1934

For the month of September 2023

Commission File Number: 001-37643

PURPLE BIOTECH LTD.
(Translation of registrant's name into English)

4 Oppenheimer Street, Science Park, Rehovot 7670104, Israel
(Address of principal executive offices)

Indicate by check mark whether the registrant files or will file annual reports under cover Form 20-F or Form 40-F.

Form 20-F Form 40-F

Purple Biotech Ltd. (the "Company" or the "Registrant") is announcing that it has made available an updated Company Presentation on its website. A copy of the updated Company Presentation is attached hereto as Exhibit 99.1 and may be viewed at the Company's website at www.purple-biotech.com.

Exhibit

99.1 [Purple Biotech Corporate Presentation September 2023](#)

Incorporation by Reference

This Report on Form 6-K, including all exhibits attached hereto, is hereby incorporated by reference into each of the Registrant's Registration Statement on [Form S-8](#) filed with the Securities and Exchange Commission on May 20, 2016 (Registration file number 333-211478), the Registrant's Registration Statement on [Form S-8](#) filed with the Securities and Exchange Commission on June 6, 2017 (Registration file number 333-218538), the Registrant's Registration Statement on [Form F-3](#), as amended, originally filed with the Securities and Exchange Commission on July 16, 2018 (Registration file number 333-226195), the Registrant's Registration Statement on [Form S-8](#) filed with the Securities and Exchange Commission on March 28, 2019 (Registration file number 333-230584), the Registrant's Registration Statement on [Form F-3](#) filed with the Securities and Exchange Commission on September 16, 2019 (Registration file number 333-233795), the Registrant's Registration Statement on [Form F-3](#) filed with the Securities and Exchange Commission on December 2, 2019 (Registration file number 333-235327), the Registrant's Registration Statement on [Form F-1](#) filed with the Securities and Exchange Commission on December 27, 2019 (Registration file number 333-235729), the Registrant's Registration Statement on [Form F-3](#) filed with the Securities and Exchange Commission on May 13, 2020 (Registration file number 333-238229), the Registrant's Registration Statement on [Form S-8](#) filed with the Securities and Exchange Commission on May 18, 2020 (Registration file number 333-238481), each of the Registrant's Registration Statements on Form F-3 filed with the Securities and Exchange Commission on July 10, 2020 (Registration file numbers [333-239807](#) and [333-233793](#)), the Registrant's Registration Statement on [Form S-8](#) filed with the Securities and Exchange Commission on April 4, 2022 (Registration file number 333-264107) and the Registrant's Registration Statement on [Form F-3](#) filed with the Securities and Exchange Commission on March 23, 2023 (Registration file number 333-270769) and the Registrant's Registration Statement on [Form F-3](#), as amended, originally filed with the Securities and Exchange Commission on December 8, 2022 (Registration file number 333-268710), to be a part thereof from the date on which this report is submitted, to the extent not superseded by documents or reports subsequently filed or furnished.

SIGNATURES

Pursuant to the requirements of the Securities Exchange Act of 1934, the Registrant has duly caused this report to be signed on its behalf by the undersigned, thereunto duly authorized.

September 13, 2023

PURPLE BIOTECH LTD.

By: /s/ Lior Fhima

Lior Fhima
Chief Financial Officer



CORPORATE PRESENTATION



NASDAQ/TASE: PPBT
September 2023

Forward-looking Statements and Safe Harbor

Certain statements in this press release that are forward-looking and not statements of historical fact are forward-looking statements within the meaning of the safe harbor provisions of the Private Securities Litigation Reform Act of 1995. Such forward-looking statements include, but are not limited to, statements that are not statements of historical fact, and may be identified by words such as "believe", "expect", "intend", "plan", "may", "should", "could", "might", "seek", "target", "will", "project", "forecast", "continue" or "anticipate" or their negatives or variations of these words or other comparable words or by the fact that these statements do not relate strictly to historical matters. You should not place undue reliance on these forward-looking statements, which are not guarantees of future performance. Forward-looking statements reflect our current views, expectations, beliefs or intentions with respect to future events, and are subject to a number of assumptions, involve known and unknown risks, many of which are beyond our control, as well as uncertainties and other factors that may cause our actual results, performance or achievements to be significantly different from any future results, performance or achievements expressed or implied by the forward-looking statements. Important factors that could cause or contribute to such differences include, among others, risks relating to: the expected benefits, synergies and costs of the transaction; management plans relating to the transaction; the potential future financial impact of the transaction; and any assumptions underlying any of the foregoing; the expected timing of the completion of the transaction and the parties' ability to complete the transaction; the plans, strategies and objectives of management for future operations; product development for NT219 and CM24 as well as Immunizon Ltd.'s portfolio of investigational tri-specific antibody compounds to be acquired; the process by which such early stage therapeutic candidates could potentially lead to an approved drug product is long and subject to highly significant risks, particularly with respect to a joint development collaboration; the fact that drug development and commercialization involves a lengthy and expensive process with uncertain outcomes; our ability to successfully develop and commercialize our pharmaceutical products; the expense, length, progress and results of any clinical trials; the impact of any changes in regulation and legislation that could affect the pharmaceutical industry; the difficulty in receiving the regulatory approvals necessary in order to commercialize our products; the difficulty of predicting actions of the U.S. Food and Drug Administration or any other applicable regulator of pharmaceutical products; the regulatory environment and changes in the health policies and regimes in the countries in which we operate; the uncertainty surrounding the actual market reception to our pharmaceutical products once cleared for marketing in a particular market; the introduction of competing products; patents obtained by competitors; dependence on the effectiveness of our patents and other protections for innovative products; our ability to obtain, maintain and defend issued patents; the commencement of any patent interference or infringement action against our patents, and our ability to prevail, obtain a favorable decision or recover damages in any such action; and the exposure to litigation, including patent litigation, and/or regulatory actions, and other factors that are discussed in our Annual Report on Form 20-F for the year ended December 31, 2022 and in our other filings with the U.S. Securities and Exchange Commission ("SEC"), including our cautionary discussion of risks and uncertainties under "Risk Factors" in our Registration Statements and Annual Reports. These are factors that we believe could cause our actual results to differ materially from expected results. Other factors besides those we have listed could also adversely affect us. Any forward-looking statement in this press release speaks only as of the date which it is made. We disclaim any intention or obligation to publicly update or revise any forward-looking statement or other information contained herein, whether as a result of new information, future events or otherwise, except as required by applicable law. You are advised, however, to consult any additional disclosures we make in our reports to the SEC, which are available on the SEC's website, <https://www.sec.gov>.



Corporate Highlights

Purple Biotech identifies promising first-in-class drug candidates to treat cancers with high unmet medical need

- Multiple data read-outs expected in 2023 & 2024
- Two First-in-Class clinical stage drugs
- A preclinical tri-specific immuno-engagers platform
- Lean & global operation
- Cash runway into 1H25

Purple Biotech (NASDAQ/TASE: PPBT)

As of June 30, 2023

- ADS Outstanding: 21.4 M
- Cash Balance : \$18 M

**Strong position to reach short and mid term
value creating clinical data catalysts**



Leadership Team



Eric K. Rowinsky, MD
Chairman of the Board
Former CMO at ImClone, Stemline,
Board member at Biogen Inc.



Gil Efron
Chief Executive Officer
Former Deputy CEO & CFO at
Kamada (NASDAQ:KMDA)

 KAMADA



Lior Fhima
Chief Financial Officer
Formerly at Kamada
(NASDAQ:KMDA)

 KAMADA



Michael Schickler, PhD
Head of Clinical and Regulatory Affairs
Formerly at Hoffmann-La Roche,
CEO at CureTech



PURPLE



Hadas Reuveni, PhD
VP Research & Development
Formerly at Keryx
(NASDAQ:KERX)





Fabien Sebille, PhD
Chief Business Officer
Formerly at
Debiopharm

 Debiopharm

A Pipeline Dedicated to Advancing Oncology Therapies

| Project | Target | Indications | Development Stage | | | | Value Drivers |
|---------|-------------------------------|--|-------------------|---------|----------|-----------|---|
| | | | Pre-Clinical | Phase I | Phase II | Phase III | |
| CM24* | CEACAM1 mAb | Pancreatic Cancer (+nivolumab+SoC) | | | | | <ul style="list-style-type: none"> Phase 2 interim analysis 2H23 & 1H24 Phase 2 top line results 2H24 |
| NT219 | STAT3xIRS1/2 Dual Inhibitor | Solid tumors (monotherapy) | | | | | <ul style="list-style-type: none"> Phase 1 results 2H23 Initiation of Phase 2 1H24 |
| | | Head and Neck & Colorectal Cancer (+Cetuximab) | | | | | |
| IM1240 | CD3x5T4xNKG2A Tri-specific Ab | Solid Tumors | | | | | |

*Clinical collaboration and supply agreement with:  Bristol Myers Squibb



Multiple data read-outs expected in the next 12 months

| 5



Advancing First-in-Class Oncology Therapies

CM24: an α -CEACAM1* mAb

**Lead indication: Pancreatic Ductal
Adenocarcinoma (PDAC)**

*Carcinoembryonic Antigen Cell Adhesion Molecule

CM24: a New Multi-Functional Immune Checkpoint Inhibitor

Attractive new target

- CEACAM1 is **overexpressed** on certain **tumor cells** and **infiltrating immune cells**
- CEACAM1 interacts with **CEACAM1** and **CEACAM5** and creates a **tumor-protective environment**

Demonstrated mechanism of action

- CM24 increases **T cell and NK cells cytotoxicity** against tumors
- CM24 shows benefit in combination with immuno-oncology treatments
- CM24 **blocks adhesion** of tumor cells to Neutrophil Extra cellular Traps (NETs)

Signals of clinical efficacy


- **Favorable safety profile** in monotherapy and in combination with nivolumab*
- **Partial response and stable disease** in dose escalation study with nivolumab
- Potential **biomarkers identified** such as NETs and CEACAM1 levels on TILs
- **Randomized Phase 2 initiated in Q1 2023**

Sizable market potential

- Significant unmet medical need in **pancreatic ductal adenocarcinoma (PDAC)**, most common form of pancreatic cancer
- Strong **IP position** and well **ahead of competitors**
- **Multiple opportunities** to leverage the MoA in other clinical settings




CM24 MOA | Immune Check Point Inhibitor & Anti-Metastatic Activity



Merkel et al. *J Immunol* 2002, 2006; *Immunology*, 2008; *Cancer Immunol Immunother* 2010; Ortenberg et al. *Mol Cancer Ther* 2012; Zhou, 2009; Li, 2013; Huang, 2015; Acharya N, et al. *J Immunotherapy Canc* 8:e911-22, 2020.; Rayes RF, et al. *Neutrophil Extracellular Trap-Associated CEACAM1 as a Putative Therapeutic Target to Prevent Metastatic Progression of Colon Carcinoma. J Immunol.* 2020; Gerstel, D. et al. *CEACAM1 creates a pro-angiogenic tumor microenvironment that supports tumor vessel maturation. Oncogene* 30, 4275-4289 (2011).

CEACAM1 is Over-Expressed in Pancreatic Cancer



Large Market Opportunity in Pancreatic Cancer



- Pancreatic Cancer accounts for ~60K new cases/year in the US alone; with a 5-year relative survival rate of 11.5%¹
- Immuno-oncology approaches have been limited to patients with high microsatellite instability (MSI-H) or high tumor mutational burden (TMB-H)
- 5-year overall survival rate with chemotherapy in 2nd line patients is 3%¹
- 3L standard of care regimens efficacy data: patients treated without chemotherapy: Overall Survival (OS) 2 months, OS of 3-4 months with chemotherapy
- 2L standard of care regimens efficacy data: Gemcitabine/Nab-paclitaxel³: OS 7.9 months, Progression Free Survival (PFS) 4.3 months or Nab-IRI/5FU/LV⁴: OS 6.2 months, PFS 3.1 months
- CEACAM1 expression correlates with poor prognosis in Pancreatic cancer²
- Preclinical data support significant synergy of CM24 with currently marketed immuno-oncology therapies

Combining nivolumab with CM24 in a clinical collaboration with Bristol Myers Squibb



1. American Cancer Society, *Cancer Facts & Figures 2019*, and the ACS website, <https://seer.cancer.gov/statfacts/html/pancreas.html>
2. Calinescu et al, *Journal of Immunology Research* 2018: 7169081; *Carcinoembryonic antigen-related cell adhesion molecules (CEACAM) 1, 5 and 6 as biomarkers in pancreatic cancer*, DOI:10.1371/journal.pone.0113023
3. De Jesus VHF, Camandaroba MPG, Calsavara VF, Riechelmann RP. Systematic review and meta-analysis of gemcitabine-based chemotherapy after FOLFIRINOX in advanced pancreatic cancer. *Therapeutic Advances in Medical Oncology*. 2020;12: do:10.1177/1758835920905408
4. Wang Gilam A, Hubner RA, Siveke JT, et al. NAPOLI-1 phase 3 study of liposomal irinotecan in metastatic pancreatic cancer: Final overall survival analysis and characteristics of long-term survivors. *Eur J Cancer*. 2019;108:78-87. do:10.1016/j.ejca.2018.12.007




CM24 Phase 1 Dose Escalation Results

Encouraging data in 2L/3L Pancreatic Ductal Adenocarcinoma (PDAC) patients

Study Results

14 patients were evaluable for efficacy:

- Best overall response included 1 Partial Response (PR) (PDAC) and 4 Stable Disease (SD) (3 PDAC and 1 papillary thyroid cancer (PTC))
- Pharmacokinetic analysis of CM24 shows exposure is dose-proportional across the 3 doses in this study
- Well tolerated with no Dose Limiting Toxicities (DLTs) and no grade ≥ 4 Adverse Events (AEs)
- Median Overall Survival 4.5 months (95% CI 2.0-11.1) for 11 PDAC patients



Phase 2 Combination Study Design (NCT04731467)

A study of CM24 in combination with nivolumab plus chemotherapy in PDAC patients in 2L treatment
18 centers are currently active in US, EU & Israel

Primary endpoint:

OS

Secondary endpoints:

PFS, OS rate @ 6 & 12 months, PFS rate @ 3 & 6 months, ORR


Interim analysis:

- PFS 2H23
- OS 1H24

Top line data:

- PFS 1H24
- OS 1H25

Measurement of CEACAM1 and other bio-markers is ongoing





Advancing First-in-Class Oncology Therapies

NT219: A Small Molecule Dual Inhibitor of IRS 1/2 and STAT3

Lead indication: Recurrent/Metastatic Head & Neck Cancer (SCCHN)

NT219 - A Novel Approach to Overcome Resistance to EGFRi and Beyond

Novel MOA

- NT219 is a **First-in-Class** small molecule
- Dual inhibitor of **IRS1/2** and **STAT3**, two major drivers of resistance to cancer therapies

Robust preclinical package

- **Significant efficacy** in multiple in-vivo models
- Modulation of the **tumor microenvironment**
- Uniquely suited to inhibit development of **drug resistance (EGFRi and KRASi)**

Clinical Stage

- Early signs of clinical **efficacy as single agent**
- **No dose limiting toxicity** observed to date, recommended dose for Phase 2 has not been determined yet

Broad Market Potential


- Opportunity to **establish a 2nd line standard of care**
- Multiple market upsides in **combination with multiple approved therapies**



NT219 blocks 2 critical signalling pathways at once

IRS1/2

- Scaffold proteins, mediating mitogenic, metastatic, angiogenic and anti-apoptotic signals from IGF1R, IR, IL4R and other oncogenes, overexpressed in multiple tumors
- Regulates major survival pathways such as the PI3K/AKT, MEK/ERK and WNT/β-catenin
- Activated as a feedback response to anti-cancer therapies
- IRS plays an important role in promoting a tumor-protective microenvironment, by mediating upregulation of TAMs and CAFs



STAT3

- Well-established transcription factor associated with the tumorigenic phenotype
- STAT3 is broadly hyperactivated in many cancers, promoting proliferation, survival, angiogenesis and metastasis
- STAT3 pathway is required for TGFβ-induced EMT and cancer cell migration and invasion
- STAT3 is a critical player in tumor immune evasion, suppressing immune stimulators and enhancing immunosuppressive factors



Hadas-Reuveni et al. Cancer Res 2012; 72:4383-4394; Machado-Neto et al. Clinics 2018; 73(suppl 1):566; Nasrwanji Iboik; Mazhar Ghaffari; Hadas Reuveni et al. Mol Cancer Ther 2014; 13(12): 2827-2839; Rampis et al. Oncogene 2016; 35(20):562-564; Flashman-Abramson, Reuveni Hadas, Levitzki Alexander et al. Oncogene 2016;35(25):2675-2680; Sanchez-Lopez et al. Oncogene 2016;35(20):2634-2644; Zhao C et al. Trends Pharmacol Sci 2016;37(1):47-56; Johnson, Daniel E et al. Nature reviews. Clinical oncology 2018; 15(4): 234-248; Zi Ying et al. J Cell Biochem 2018;139:9419-9432.

Strong Efficacy in Monotherapy




Animal model

Head & Neck Cancer
(SCC-9) NSG™, immuno-
competent



Drugs

α-PD1
Cetuximab
NT219



- NSG mice were injected SC with SCC-9 cells.
- PBMCs (18×10^6 cells per mouse) administered 4 weeks prior to first treatment.
- NT219 (IV), α-PD1 (IP), and cetuximab (IP) were administered twice a week for 4 weeks.

NT219 Restores Sensitivity to EGFRi in PDX Models

Lung Cancer




Non-small cell lung cancer (NSCLC)
Exon 19 deletion EGFR and T790M, biopsy of bone marrow metastasis, patient previously progressed on afatinib and osimertinib


Head & Neck Cancer



Recurrent/metastatic squamous cell carcinoma of the head and neck (R/M SCCHN)
metastasis, patient progressed on chemoradiation, several chemotherapies and pembrolizumab



Osimertinib 5 mg/kg, NT219 65 mg/kg, mean tumor volume at the end point, 3 mice/group;



Treatments on days 0, 3 and 10, cetuximab - 1mg/mouse, 3 mice/group; PBMCs (1.4M cells/mouse) were injected on day 6

** p<0.01, * p<0.02 based on one-way ANOVA with post hoc Tukey's HSD test

First Market Opportunity

Recurrent or Metastatic Squamous Cell Carcinoma of Head and Neck (SCCHN)




Targeting the unmet medical need

- SCCHN is the 6th most common cancer type ; 175k new cases/year are expected by 2024
- 1L standard of care has shifted from chemotherapy towards immuno-oncology + chemotherapy
- < 20% of R/M SCCHN patients respond to Pembrolizumab
- Market size forecasted to >\$5b in 2030

Rationale for combining Cetuximab + NT219

- EGFR and PD(L)-1 are the only clinically validated targets in SCCHN
- < 15% of R/M SCCHN patients respond to Cetuximab
- Cetuximab inhibits EGFR signaling and promotes ADCC in EGFR expressing tumors
- STAT3 and IRS-to-AKT activation contributes to resistance to cetuximab in SCCHN



NT219 + Cetuximab has potential to become SOC as 2nd line therapy r/m SCCHN



Global Data 2018: Head and Neck Squamous Cell Carcinoma: Opportunity Analysis and Forecasts to 2026 ; Internal best current estimates of patient numbers based on external research, 8 major global territories


| 18

NT219 Phase 1 Dose Escalation Monotherapy

Interim Results

Study Design

- As of data cutoff date of May 12th, 2022, a total of 14 patients were enrolled and 12 patients were evaluable for dose limiting toxicity (DLT) determination (4 colorectal cancer, 3 pancreatic cancer, 2 breast cancer, 1 GEJ, 1 esophageal, and 1 appendiceal cancer)
- Median number of prior treatment regimens for metastatic disease was 4 (2-11).



Safety


- No DLTs were observed across all dose levels.
- Nine Grade 3 adverse events (AEs) were observed, two of which possibly related to NT219

| AE Term | Total | 1 | 2 | 3 | Grade 4/5 |
|---|-------|---|---|---|-----------|
| Fatigue | 6 | 6 | | | |
| Constipation | 4 | 4 | | | |
| ALP increased | 3 | 2 | | 1 | |
| ALT increased | 3 | 1 | 2 | | |
| Anemia | 3 | 1 | 2 | | |
| AST increased | 3 | 1 | 1 | 1 | |
| Diarrhea | 3 | 2 | | 1 | |
| Headache | 3 | 3 | | | |
| Nausea | 3 | 2 | 1 | | |
| Abdominal pain | 2 | 1 | 1 | | |
| Belching | 2 | 2 | | | |
| Cough | 2 | 2 | | | |
| Dizziness | 2 | 2 | | | |
| Dyspnea | 2 | 2 | | | |
| Edema limbs | 2 | 2 | | | |
| Fever | 2 | 2 | | | |
| Hot flashes | 2 | 2 | | | |
| Hyperhidrosis | 2 | 2 | | | |
| Urinary tract infection | 2 | | 2 | | |
| Closed displaced fracture of right femoral neck | 1 | | | 1 | |
| Intractable right hip pain | 1 | | | 1 | |
| Malignant hypercalcemia | 1 | | | 1 | |
| Toxic Encephalopathy | 1 | | | 1 | |
| Worsening back pain | 1 | | | 1 | |
| Abdominopelvic Ascites | 1 | | | 1 | |

NT219 Phase 1 Dose Escalation Monotherapy

Interim Results: Encouraging Initial Efficacy Signals

- For the 12 evaluable patients, best overall response included **one confirmed Partial Response** (gastroesophageal junction (GEJ) patient, > 5.5 months duration of response following end of treatment), and **3 Stable Disease** with one patient awaiting follow up MRI/CT scans
- As of the cutoff date (May 12th , 2022), **10/12** patients are either **on treatment or in follow up** (range 1.1 to 18 months).



Durable Partial Response in a GEJ patient and Stable Disease in 3 out of 4 mutated KRAS colorectal cancer patients



NT219 Monotherapy and Combination Phase 1/2 Study Design (NCT04474470)

Study Title

A Phase 1/2 study with open-label, dose escalation phase followed by single-arm expansion to assess the safety, tolerability, PK, PD and efficacy of NT219, alone in adults with recurrent or metastatic solid tumors and in combination with Erbitux® (cetuximab) in Head and Neck cancer

Endpoints


Primary endpoints:

Safety, pharmacokinetics and to determine the maximum tolerated dose (MTD)

Secondary endpoints:

Obtain preliminary efficacy data


Study Design



*Colorectal Adenocarcinoma pts will be recruited in the Dose Escalation phase

■■■ = Indication TBD (expansion not part of the study protocol)

Dose Escalation Design



- Recruitment for the above dose levels is completed
- No DLTs, no significant safety concerns have been observed



Advancing First-in-Class Oncology Therapies

**IM1240: CD3x5T4xNKG2A
Conditionally-Activated Tri-Specific Antibody**

A Novel Mechanism of Action Tri-Specific Antibody

- **Multi-specific biologics is an expanding class of drugs getting a lot of interest in the industry**
- After initial success in hemato-oncology, **new formats are being investigated in solid tumors**
- Technology displays **several distinctive features:**
 - Dual engagement of T cells and NK cells to mount an optimal anti-tumoral immune response
 - A tumor-restricted activation through a **cleavable capping system** designed to provide a wide therapeutic index
 - Carefully selected Tumor Associated Antigens allowing **patient-centric development**




Removal of T cell engager's masking by
proteases of the tumor microenvironment

T and NK Cell activation
restricted to selected tumors


5T4, a Novel Target in Oncology

5T4 is highly expressed on certain tumors and correlates with poor prognosis



Am J Cancer Res 2018;8(4):610-623 www.ajcr.us /ISSN:2156-6976/ajcr0074519


5T4 is a Tumor Associated Antigen prevalent to several large indications




Opportunity of patient stratification strategy (5T4⁺)

| 24

Promising Proof of Concept Data



- Tribody induces **sub-nM EC₅₀** against A549 cells
- **20-fold more potent** than CD3x5T4 variant
- Cell killing validated on **multiple 5T4⁺ cell lines** (MDA-MB-231, HCT116, NCI-H226)



- **Sustained regressions** in Breast Cancer xenograft model (MDA-MB-231)
- The Pro-Tribody Capped-CD3x5T4xNKG2A performed better than the uncapped variant
- **No change** in body weight



Corporate Highlights

Purple Biotech identifies promising first-in-class drug candidates to treat cancers with high unmet medical need

- Multiple data read-outs expected in 2023 & 2024
- Two First-in-Class clinical stage drugs
- A preclinical tri-specific immuno-engagers platform
- Lean & global operation
- Cash runway into 1H25

Purple Biotech (NASDAQ/TASE: PPBT)

As of June 30, 2023

- ADS Outstanding: 21.4 M
- Cash Balance : \$18 M

**Strong position to reach short and mid term
value creating clinical data catalysts**





PURPLE
BIOTECH

THANK YOU





Contact Us:
ir@purple-biotech.com



Appendix A | CM24

CM24 Reduces Tumor Burden & Synergetic with α -PD-1




Significant benefits as both single agent and in combination with α -PD-1

Phase 1 Dose Escalation Interim Results

CM24 is Safe and Well Tolerated in Combination with Nivolumab

Study Design

- As of March 8th, 2022, a total of 13 patients were enrolled and 11 patients were evaluable for DLT determination (8 PDAC, 2 CRC and 1 PTC).
- 9 patients had received 2 prior regimens for metastatic disease, 2 patients had one previous line.



Safety

- No DLTs were observed across all dose levels; no Grade 4 AEs or treatment-related deaths have been reported.
- Grade 3 AEs were noted in 6/13 patients (46%).

| AE Term | Total | Grade | | | |
|-------------------------------|-------|-------|---|---|-----|
| | | 1 | 2 | 3 | 4/5 |
| Diarrhea | 5 | 4 | | 1 | |
| Abdominal pain | 4 | 1 | 3 | | |
| Fever | 4 | 2 | 2 | | |
| Headache | 4 | 3 | 1 | | |
| Fatigue | 4 | 4 | | | |
| Nausea | 3 | 1 | 2 | | |
| Creatinine increased | 3 | 2 | 1 | | |
| Hypokalemia | 2 | | | 2 | |
| Dyspnea | 2 | 1 | | 1 | |
| Constipation | 2 | 2 | | | |
| Cough | 2 | 2 | | | |
| Abdominal pain aggravated | 1 | | | 1 | |
| Alkaline phosphatase increase | 1 | | | 1 | |
| Atrial flutter | 1 | | | 1 | |
| C-Diff Colitis | 1 | | | 1 | |
| GI bleed | 1 | | | 1 | |
| Leukocytosis | 1 | | | 1 | |
| Small bowel obstruction | 1 | | | 1 | |



CM24 Phase 1 Combination Study (NCT04731467)

Demographics

In the Phase 1 part, patients with indicated refractory cancers were administered CM24 at 10, 15, and 20mg/kg q2w and nivolumab 480mg q4w.

- The primary objective of this part was to evaluate safety, tolerability, pharmacokinetics and determine the RP2D
- Safety was assessed according to CTCAE v5 and preliminary anti-tumor activity was assessed by the investigators according to RECISTv1.1 using CT/MRI
- CM24 and CEACAM1 measurements in serum, biopsy specimens, and TILs, as well as tumor and TILs PD-L1 levels are being determined

As of March 8th, 2022, a total of 13 patients were enrolled and 11 patients were evaluable for dose-limiting toxicity (DLT) determination (8 PDAC, 2 CRC and 1 PTC)

- 9 patients had received 2 prior regimens for metastatic disease and 2 patients had one previous line.

Demographics of patients treated with CM24 (10, 15, 20mg/kg) in combination with nivolumab (480mg)


| Median age, years (range) | 65 (49-76) | Prior Lines of Therapy, n (%) | |
|---------------------------|------------|---|------------|
| Sex, n (%) | | 1 | 2 (18%) |
| Male | 5 (45%) | 2 | 9 (82%) |
| Female | 6 (55%) | Diagnosis, n (%) | |
| Ethnicity, n (%) | | Pancreatic cancer | 8 (73%) |
| Not Hispanic or Latino | 10 (91%) | Papillary Thyroid cancer | 1 (9%) |
| Hispanic or Latino | 1 (9%) | Colorectal cancer | 2 (18%) |
| Race, n (%) | | Median Time from Initial Diagnosis months (range) | 23 (11-73) |
| White | 10 (91%) | ECOG, n (%) | |
| Black or African American | 1 (9%) | 0 | 7 (64%) |
| | | 1 | 4 (36%) |



Confirmed Partial Response in a 3L PDAC Patient

Patient Profile

- 65 y/o female, pancreatic cancer
- 2 prior lines of treatments: FOLFIRINOX and gemcitabine/nab-paclitaxel
- Post Whipple Procedure
- Patient had a germline NF1 VUS, with MSI-S and PDL-1 IHC 2+ and 5% staining
- Confirmed Partial Response: after initial treatment, the patient had a Partial Response of 40%, with a definite reduction of the para-tracheal adenopathy and liver lesions and 58% reduction in CA19-9 levels
- Under treatment for 6 months, still under monitoring.



CEACAM1 Plays a Key Role in Cancer Biology

01 | ADHESION

Horst, 2011

Oncogene

"CEACAM1 creates a pro-angiogenic tumor microenvironment that supports tumor vessel maturation"

Ferrari, 2020

 *Journal of Immunology*

"Neutrophil extracellular trap-associated CEACAM1 as a putative therapeutic target to prevent metastatic progression of colon carcinoma"



02 | IMMUNE CELLS/ IMMUNE EXCLUSION

Tsuzuki, 2020



"Immune-checkpoint molecules on regulatory T-cells as a potential therapeutic target in head and neck squamous cell cancers"

Tsang, 2020



"[Blockade] enhances natural killer cell cytotoxicity against tumor cells through blockade of the inhibitory CEACAM1 / CEACAM5 immune checkpoint pathway"


03 | IMMUNO-ONCOLOGY

Blumberg, 2015

nature

"CEACAM1 regulates TIM-3-mediated tolerance and exhaustion"

Shively, 2013




"CEACAM1 regulates Fas-mediated apoptosis in Jurkat T-cells via its interaction with β -catenin"



Appendix B | NT219

Novel MOA: IRS Degradation By NT219 Blocking IGF1R-AKT Pathway¹



¹Reuveni et al. *Cancer Res* 2013 ; Ibuki et al. *Mol Cancer Ther* 2014



NT219 Re-sensitizes Tumors Refractory to α -PD1




PDX Model

Humanized PDX of
GastroEsophageal Junction
(GEJ) Cancer (refractory to
pembrolizumab)



Drug

Pembrolizumab
(Keytruda®)




* Double autologous model - Tumors & PBMCs are from the same patient (#RA236) | Keytruda - 6mg/kg IP, NT219 - 60mg/kg IV

Novel MOA

Signal Transducer and Activator of Transcription 3 (STAT3) Inhibition

- Point of convergence for numerous oncogenic signaling pathways
- Central in regulating the anti-tumor immune response
- Broadly hyperactivated both in cancer and non-cancerous cells within the tumor ecosystem and plays important roles in inhibiting the expression of crucial immune activation regulators and promoting the production of immunosuppressive factors
- Targeting the STAT3 signaling pathway has emerged as a promising therapeutic strategy for numerous cancers


NT219 demonstrates a durable and dose-dependent suppression of STAT3 tyrosine phosphorylation, affecting both the tumor cells and the tumor microenvironment.





Zou, S., Tong, Q., Liu, B., et al. Targeting STAT3 in Cancer Immunotherapy. *Mol Cancer* 19, 145 (2020). <https://doi.org/10.1186/s12943-020-01258-7>

Simultaneous Blockade of STAT3 and AKT Pathways are Required to Overcome Resistance to EGFRi

Overcoming drug resistance



Proof of Concept: PDX model of Head and Neck Cancer



By blocking both STAT3 and IRS resistance pathways, NT219 re-sensitizes tumors to anti-cancer therapies



NT219 Phase 1 Dose Escalation Monotherapy Demographics

As of data cutoff date of May 12th , 2022, a total of 14 patients were enrolled to 4 NT219 dose levels (3 - 24mg/kg)

- 12 patients were evaluable for DLT determination including 4 CRC, 3 pancreatic cancer, 2 breast cancer, and one of each of the following cancers: GEJ, esophageal and appendiceal cancer
- Median number of prior treatment regimens for metastatic disease was 4 (2-11)

| Demographics of patients treated with NT219 (3, 6, 12, 24mg/kg) | | | |
|---|------------|-------------------------------|---------|
| Median age, years (range) | 67 (39-79) | Diagnosis , n (%) | |
| Sex, n (%) | | Pancreatic | 3(25%) |
| Male | 4(33%) | GE Junction | 1(8%) |
| Female | 8 (67%) | Breast | 2(17%) |
| Ethnicity, n (%) | | Colorectal | 4(33%) |
| Not Hispanic or Latino | 11 (92%) | Appendiceal | 1(8%) |
| Hispanic or Latino | 1 (8%) | SCC of the esophagus | 1(8%) |
| Race, n (%) | | Prior Lines of Therapy, n (%) | |
| White | 10 (83%) | 2 | 2 (17%) |
| Black or African American | 2 (17%) | 3 | 3 (25%) |
| ECOG, n (%) | | 4 | 2(17%) |
| 0 | 5 (42%) | 5 | 2(17%) |
| 1 | 7 (58%) | 6 | 1(8%) |
| Median Time from Initial Diagnosis months (range) | 36(10-153) | 8 | 1(8%) |
| | | 11 | 1(8%) |



Selected Publications



Michael
Karin

Oncogene (2012) 31, 2639–2648
© 2012 Macmillan Publishers Limited. All rights reserved 0950-2126/12
www.nature.com/onc

ORIGINAL ARTICLE

Targeting colorectal cancer via its microenvironment by inhibiting IGF-1 receptor-insulin receptor substrate and STAT3 signaling

E Sanchez-Lopez¹, E Flaherman-Alberman², S Shulman¹, Z Zhong¹, K Tengs^{1,3}, A Levitzki¹ and M Karin¹



Alexander
Levitzki

Oncogene (2012) 31, 2649–2656
© 2012 Macmillan Publishers Limited. All rights reserved 0950-2126/12
www.nature.com/onc

SHORT COMMUNICATION

Targeting melanoma with NT157 by blocking Stat3 and IGF1R signaling

E Flaherman-Alberman¹, S Kim¹, C Huh¹, E Shulman¹, H Song¹, A Shai¹, Y Leng¹, M Bar Eli¹, H Assouline^{1,2,3} and A Levitzki¹

THE UNIVERSITY OF TEXAS
MD Anderson
Cancer Center

Menashe
Bar-Eli

Published OnlineFirst May 7, 2013; DOI: 10.1158/0008-5472.CAN-12-3385

Therapeutics, Targets, and Chemical Biology



Therapeutic Destruction of Insulin Receptor Substrates for Cancer Treatment

Hadas Reuveni^{1,2}, Eyal Flaherman-Alberman¹, Uteck Shiner^{1,2}, Kati Makarowski^{1,2}, Hendrik Song¹,
Assaf Shai¹, Meiriont Heny¹, Menashe Bar-Eli¹, and Alexander Levitzki¹



Michael
Cox

Published OnlineFirst September 29, 2014; DOI: 10.1158/1538-7463.MCT-13-0842

Small Molecule Therapeutics




The Tyrosostatin NT157 Suppresses Insulin Receptor Substrates and Augments Therapeutic Response of Prostate Cancer

Henkaku Itoh^{1,2}, Mazene Ghofer^{1,2}, Hadas Reuveni^{1,2}, Mital Pandey¹, Ladan Fadi¹, Hanan Assouline¹,
Martin E. Gleisner^{1,3}, Alexander Levitzki¹, and Michael E. Cox^{1,4}



NT219 Suppresses β -Catenin activity in CRC Cells and Inhibited CRC Brain Metastasis



Colon cancer LS-513 cells overexpressing IRS2 demonstrate enhanced β -catenin activity.

Targeted inhibition of IRS2 by NT219 or IRS2-SH RNA, suppresses the increased β -catenin activity and inhibit LS-513 cell viability.

Combination of 5-FU and NT219 significantly inhibited the growth of CRC tumors in brain, using intracranial model and extended mice survival.



AACR Annual Meeting, April 2021, AACR Virtual Special Conference on Epigenetics and Metabolism, Oct 2020, Ido Wolf, MD, Head of Oncology Division, Tel Aviv Sourasky Medical Center

NT219 | Pancreatic Cancer in Combination with Gemcitabine



PDX model

Pancreatic Cancer







Drug

Gemcitabine (Gemzar®)



PURPLE

Highly effective anti cancer activity exhibited by NT219 in combination with Gemcitabine



RNA Sequencing | Analysis of Tumors Following Treatment



PDX model

Pancreatic Cancer





Drug

Gemcitabine (Gemzar®)




Reduced expression of IRS1, Ki67, FOXM1 & TGFb is
exhibited by pancreatic cancer treated with NT219 alone and
in combination with gemcitabine



Confirmed Partial Response as a Single Agent in a GEJ Cancer Patient

- In a patient with refractory GE junction disease (mutated KRAS, TP53), NT219 administration (3mg/kg as a single agent) was associated with a confirmed partial response (PR):
 - Complete remission at the largest target lesion (right)
 - Complete resolution of all non-target lesions (two lymph nodes) has also been demonstrated
 - The patient remained on treatment for nearly 6 months.



GEJ tumor at baseline screening

CT imaging of the GEJ tumor
after 5 months of treatment with
NT219

